



Nitrogen-corrected True Metabolizable Energy and Amino Acid Digestibility of Chinese Corn Distillers Dried Grains with Solubles in Adult Cecectomized Roosters

F. Li^a, Y. Liu^{1,a}, R. Q. Yin, X. J. Yang, J. H. Yao*, F. F. Sun, G. J. Li, Y. R. Liu¹ and Y. J. Sun¹
College of Animal Science and Technology, Northwest A&F University, Yangling, Shaanxi 712100, China

ABSTRACT: This study was conducted to evaluate chemical composition, nitrogen-corrected true metabolizable energy (TME_n) and true amino acids digestibility of corn distillers dried grains with solubles (DDGS) produced in China. Twenty five sources of corn DDGS was collected from 8 provinces of China. A precision-fed rooster assay was used to determine TME_n and amino acids digestibility with 35 adult cecectomized roosters, in which each DDGS sample was tube fed (30 g). The average content of ash, crude protein, total amino acid, ether extract, crude fiber and neutral detergent fiber were 4.81, 27.91, 22.51, 15.22, 6.35 and 37.58%, respectively. TME_n of DDGS ranged from 1,779 to 3,071 kcal/kg and averaged 2,517 kcal/kg. Coefficient of variation for non-amino acid crude protein, ether extract, crude fiber and TME_n were 55.0, 15.7, 15.9 and 17.1%, respectively. The average true amino acid digestibility was 77.32%. Stepwise regression analysis obtained the following equation: TME_n, kcal/kg = -2,995.6+0.88×gross energy+49.63×a* (BIC = 248.8; RMSE = 190.8; p<0.01). Removing gross energy from the model obtained the following equation: TME_n, kcal/kg = 57.88×ether extracts+87.62×a* (BIC = 254.3, RMSE = 223.5; p<0.01). No correlation was found between color scores and lysine true digestibility (p>0.05). These results suggest that corn DDGS produced in China has a large variation in chemical composition, and gross energy and a* value can be used to generate TME_n predict equation. (**Key Words:** Amino Acid, Corn Distillers Dried Grains with Solubles, Nitrogen-corrected True Metabolizable Energy, Rooster)

INTRODUCTION

Distillers dried grains with solubles (DDGS) has been commonly used as an ingredient in poultry diets for many decades. A recent study indicated that diets containing up to 24% DDGS had no adverse effect on performance of broilers (Shim et al., 2011). However, large variability of DDGS composition and poor protein quality may ultimately limit its utilization in poultry diets (Cromwell et al., 1993; Fastinger et al., 2006). Many studies were conducted to develop some useful parameters (i.e. color scores or chemical composition) to predict amino acid digestibility and nitrogen-corrected true metabolizable energy (TME_n) of DDGS (Batal and Dale, 2006; Fastinger et al., 2006). Color score was found to be a fast method to determine corn

DDGS quality (Ergul et al., 2003). A previous study has found that lighter colored DDGS had a greater concentration of bioavailable lysine (Pahm et al., 2009). In addition, some regression equations have been established to predict energy utilization based on chemical composition for poultry. Several equations were established to estimate the TME_n based on proximate composition, however, the accuracy of the prediction equation was low (Batal and Dale, 2006). And the color scores included in the model also improved the equation for wheat DDGS (Cozannet et al., 2010).

Unlike commercial databases, China DDGS nutrient data were always reported as the average value of many samples and lacked of proper prediction methods. This results in some misleading conception in using DDGS in poultry diets. The objective of this study was to determine the chemical composition and amino acids (AA) digestibility of Chinese corn DDGS and their correlation with color scores. The second objective was to establish a TME_n predicting equation for cecectomized roosters based on chemical composition and color scores.

* Corresponding Author: J. H. Yao. Tel: +86-29-87092102, Fax: +86-29-87092164, E-mail: yaojunhu2004@sohu.com

¹ Shenzhen Kondarl Gaoling Feed Co. Ltd., Gaoling Shaanxi, 710200. China.

^a These authors contributed equally to this paper.

Submitted Oct. 25, 2012; Accepted Dec. 31, 2012; Revised Jan. 28, 2013

MATERIAL AND METHODS

Samples, animals and experimental design

All procedures used in this study were approved by the Animal Care Committee of Northwest A&F University.

Twenty five corn DDGS samples (1 to 2 kg per sample) were collected from 8 provinces of China. Samples were ground through a 1 mm screen and stored at -20°C for further analysis. Thirty-five roosters were cecectomized according to the procedures described as Parsons (1985) at the age of 45 wk. After 1 month recovery, the roosters were randomly divided into 7 groups, 5 birds per group with similar body weight. Roosters were individually housed in stainless cages.

An experiment with 6 periods was used to determine TME_n and true digestible amino acid (TDAA) of DDGS. From periods 1 to period 3, seven DDGS samples were randomly assigned to each group with 5 replicates (1 bird per replicate). The other four samples were studied in period 4 with 5 replicates (1 bird per replicate) per group. In each period, birds were given 50 ml of 60% glucose solution after a 24 h fast. After a further 24 h, each bird was accurately tube fed 30 g of DDGS. All excreta were collected for the subsequent 48 h. Excreta samples were weighted, 10% HCl (1 ml/10 g excreta) added and stored at -20°C. After sample collection, all birds were fed a conventional corn-soybean meal diet for approximately 10 to 14 d to recovery body condition and randomly assigned to each group based on similar body weight before next period. Another 2 periods were used to estimate individual bird endogenous AA and energy losses. All roosters were fasted for 24 h and then perfused with a glucose solution. The excreta were collected for 48 h as described.

Samples analyses

Color scores including degree of lightness (L^*), redness (a^*), and yellowness (b^*) for the DDGS samples were measured using a color-difference meter (WSC-S, Shanghaishengguang Co., Ltd., Shanghai, China). Samples were held tight and flat in the test container. Reported color scores were the mean of 5 measurements, with the sample being re-mixed and leveled between each determination. Low values for L^* indicated a dark color; whereas higher scores indicated a light color (0 = black; 100 = white). Higher values of a^* and b^* indicated greater degrees of redness and yellowness, respectively.

All samples were analyzed for proximate composition DM, CP, ether extracts (EE), crude fiber (CF), ash and acid detergent fiber (ADF) by the method of AOAC (2006). Neutral detergent fiber (NDF) was determined according to Van Soest et al. (1991). Amino acid in DDGS samples and excreta were determined using an amino acid analyzer (Model 121MB, Beckman Coulter, Inc., Fullerton, CA,

USA) with the method outlined by AOAC methods 994.12 (sulfur and regular). Gross energy (GE) was analyzed using an adiabatic oxygen bomb calorimeter (Model 6300, Parr Instruments, Moline, IL, USA). Reported values were expressed as the average of duplicate sample.

Calculations

True digestibility of AA (TDAA) using the following equation:

$$TDAA (\%) = ((AA \text{ intake} + \text{endogenous AA loss} - \text{excreta AA}) / AA \text{ intake}) \times 100.$$

TME_n was calculated according to the procedure outlined by Farhat et al. (1998).

$$TME_n = (GI - GO) / FI + (FGL / FI) - (8.22 \times ANR / FI) - (8.22 \times FNL / FI);$$

where GI is the gross energy intake (kcal), GO is gross energy output (kcal), FI is feed intake of the DDGS (30 g), FGL is the fasting gross energy loss (kcal) from the feed deprived birds, ANR is apparent nitrogen retention (g), and FNL is fasting N loss (g). The gross energy excreted was corrected to zero-N balance using a factor of 8.22 kcal/g. Non-amino acids CP (NAACP) was in DDGS samples were calculated by CP minus total AA.

Statistical analysis

Coefficient of variation (CV) was calculated among all DDGS samples. Pearson's correlation analysis was performed to establish the relationship between chemical compositions and color score (using PROC CORR; SAS Institute Inc., Cary, NC, 2003). Significance was declared at $p < 0.05$.

Outliers were determined when the values (ratio of TME_n to GE) were more than 2 SD above or below the mean. Two samples were removed from the data set. The regression analysis was performed by the new database. Prediction equations of TME_n with individual proximate compositions and color scores were established by regression analysis with stepwise method (using PROC REG of SAS). The best fit equations were defined by Root MSE, Mallows statistic $C(p)$ and BIC. If the intercept was not significant ($p > 0.15$) in the equation, it was removed from the model and the R^2 was calculated using the NOINT option of SAS.

RESULTS

Nutrient composition and color score

Nutrient composition and color scores of 25 DDGS samples are presented in Table 1. The GE ranged from

Table 1. Nutrient composition and color scores of corn distillers dried grains with solubles (as-fed bases, n = 25)

Items ¹	Maximum	Minimum	Means±SD	CV (%)
Gross energy (kcal/kg)	5,164.6	4,214.7	4,642.0±227.2	4.9
TME _n (kcal/kg)	2,781.7	1,171.7	2,143.8±366.0	17.1
Dry matter (%)	91.11	84.18	87.44±1.92	2.2
Ash (%)	7.45	3.37	4.81±0.87	18.1
Crude protein (%)	34.44	23.96	27.91±2.70	9.7
Crude fiber (%)	10.04	5.12	6.35±1.01	15.9
Ether extract (%)	19.42	9.82	15.22±2.39	15.7
Neutral detergent fiber (%)	48.71	27.04	37.58±3.65	9.7
Acid detergent fiber (%)	14.85	3.67	10.61±2.49	23.5
Nitrogen-free extract (%)	39.15	25.17	33.14±3.71	11.2
Total amino acid (%)	25.52	17.15	22.51±1.93	8.6
NAACP (%)	11.82	0.14	5.40±2.97	55.0
L*(Lightness)	58.27	36.95	47.00±5.87	12.5
a*(Redness)	22.03	14.45	18.07±1.64	9.1
b*(Yellowness)	44.22	26.22	36.27±4.82	13.3

¹ NAACP = Non-amino acid crude protein, SD = Standard deviation, CV = Coefficient of variation.

4,215 to 5,147 kcal/kg and average 4,642 kcal/kg. Data of TME_n ranged from 1,172 to 2,782 kcal/kg and 17.1% of CV. Among all proximate compositions, ADF had the highest variation (CV = 23.5%), and DM had the lowest variation (CV = 2.2%). CV for the other nutrients ranged from 4.9% to 15.9%. EE content ranged from 9.80 to 19.40% and averaged 15.22%. Values of NAACP between 0.14 and 11.82% and 55.0% of CV. CV for L*, a* and b* were 12.5%, 9.1% and 13.3%, respectively.

Among all DDGS sources (Table 2), glutamic acid was the highest (3.69%), followed by leucine (2.26%), proline (1.74%) and alanine (1.71%), while cystine concentration

was the lowest (0.32%). The highest TDAA was arginine (86.79%), followed by alanine (83.88%) and leucine (83.47%). True digestible methionine and lysine were 75.28% and 80.86%, respectively.

Correlation between variables

Pearson's correlation coefficients among nutrient composition and color scores (Table 3) demonstrated that TME_n was positively correlated with GE and EE (p<0.05) and negatively correlated with CF (p<0.01). GE was (p<0.05) positively correlated with EE and color scores. There were no significant (p>0.10) correlations of color

Table 2. Amino acid content and digestibility of different corn distillers dried grains with solubles sources (as feed bases, n = 25)

Amino acids	Amino acid content (%)			TDAA ¹ (%)		
	Maximum	Minimum	Means±SD	Maximum	Minimum	Means±SD
Asp	1.96	1.12	1.57±0.18	83.07	54.72	72.43±7.40
Thr	1.39	0.21	0.87±0.09	82.94	48.48	68.78±9.04
Ser	2.79	0.70	1.15±0.36	91.93	67.90	79.13±5.74
Glu	4.35	0.88	3.69±0.69	88.46	64.30	80.92±8.09
Pro	2.38	0.58	1.74±0.37	91.20	58.47	79.92±7.15
Ala	3.58	0.44	1.71±0.48	92.73	72.78	83.88±4.37
Cys	0.44	0.14	0.32±0.08	94.33	42.94	75.83±6.96
Val	2.39	0.95	1.46±0.44	90.28	63.38	78.96±6.92
Met	0.52	0.21	0.38±0.09	95.61	44.33	75.28±11.50
Ile	1.03	0.50	0.76±0.12	83.55	49.65	73.85±7.73
Leu	2.79	1.51	2.26±0.30	89.07	74.68	83.47±3.61
Tyr	1.01	0.46	0.73±0.14	89.23	61.35	80.56±6.88
Phe	1.39	0.21	1.03±0.31	93.54	45.13	80.99±9.62
Lys	0.97	0.50	0.72±0.12	74.70	46.29	80.86±5.50
His	0.82	0.50	0.71±0.07	87.67	66.70	79.76±5.68
Arg	1.51	0.75	1.00±0.16	95.50	75.66	86.79±4.93

¹ TDAA = True digestible amino acid, SD = Standard deviation.

Table 3. Correlation matrix of TME_n with nutrient composition and color scores of corn distillers dried grains with solubles (DM basis, n = 25)

Items ¹	TME _n	GE	Ash	CP	CF	EE	NDF	ADF	NFE	L*	a*	b*
TME _n	1.00											
GE	0.46**	1.00										
Ash	0.07	-0.22	1.00									
CP	-0.29	0.12	0.09	1.00								
CF	-0.35	-0.44*	0.07	0.09	1.00							
EE	0.40*	0.67*	-0.14	-0.15	-0.23	1.00						
NDF	-0.22	-0.14	0.20	0.28	0.52**	-0.06	1.00					
ADF	-0.13	-0.36	-0.04	-0.27	0.53**	-0.31	0.13	1.00				
NFE	0.06	-0.34	-0.25	-0.75**	-0.23	-0.42*	-0.39	0.27	1.00			
L*	0.20	0.45*	0.11	0.07	0.05	0.29	0.07	0.10	-0.29	1.00		
a*	0.23	0.52**	-0.07	0.25	-0.26	0.23	-0.25	0.01	-0.25	0.59**	1.00	
b*	0.36	0.72**	-0.08	0.10	-0.29	0.49*	-0.14	-0.07	-0.29	0.84**	0.80**	1.00

GE = Gross energy, CP = Crude protein, CF = Crude fiber, EE = Ether extract, NDF = Neutral detergent fiber, ADF = Acid detergent fiber, NFE = Nitrogen-free extract.

* p<0.05; ** p<0.01.

scores with ash, CP, CF, NDF, ADF and NFE. Ether extract was positively correlated with b* value (p<0.05) and negatively (p<0.05) correlated with NFE.

Correlation relationship of crude protein and amino acid digestibility parameters with color scores are presented in Table 4. NAACP was positively correlated with CP and negatively correlated with TAA (p<0.05).

Prediction equations

Stepwise selection method was used to estimate the GE, chemical composition and color score that contributed to the predictors of TME_n of 23 DDGS samples (Table 5). GE was the first independent variable included in the model in equation 1 ($R^2 = 0.65$; BIC = 249.3; RMSE = 199.3). The model was improved with the addition a* in equation 2 with a R^2 of 0.69. And the best fit equation was: TME_n, kcal/kg = $-2,995.6 + 0.88 \times GE + 49.63 \times a^*$ ($R^2 = 0.69$; BIC = 248.8; RMSE = 190.8) at DM bases. After removing GE from the model, the best single factor to predict TME_n was b* value

(equation 3; RMSE = 259.5), followed by EE and a*. When a* was included in the model, b* was discarded (p = 0.69). In this condition, the best fit model was: TME_n, kcal/kg = $57.88 \times EE + 87.62 \times a^*$ (BIC = 254.3, RMSE = 223.5) at DM bases.

DISCUSSION

In general, excepting DM and GE, nutrient composition in the 25 sources of DDGS produced in China had considerable differences in the reported data. The average concentration of DM, CP, CF and ADF of DDGS in the present study are in agreement with the published values (Batal and Dale, 2006; Pedersen et al., 2007; Waldroup et al., 2007), while content of ash and NDF are higher than the results reported by Pedersen (2007), but similar with the results of Stein et al. (2006) (n = 10, NDF = 40.1%). Values of EE are higher than the values (9.00, 9.69 and 9.88%, respectively) reported by NRC (1994), Spiels et al. (2002)

Table 4. Correlation matrix of CP with related parameters and color scores of corn distillers dried grains with solubles (DM basis, n = 25)

Item ¹	CP	TAA	NAACP	TDCP	TDAA	TD lysine	Lysine
CP	1.00						
TAA	0.21	1.00					
NAACP	0.78**	-0.46*	1.00				
TDCP	-0.06	-0.02	-0.04	1.00			
TDAA	-0.28	-0.10	-0.19	-0.08	1.00		
TD lysine	-0.28	-0.05	-0.22	0.29	0.42**	1.00	
Lysine	0.08	0.08	0.02	0.42**	-0.31	0.39	1.00
L*	0.09	0.11	0.01	-0.18	-0.03	-0.21	0.09
a*	0.21	0.03	0.17	-0.37	-0.03	-0.26	-0.22
b*	0.10	0.01	0.09	-0.19	-0.06	-0.33	-0.09

¹TAA = Total amino acid, NAACP = Non-amino acid crude protein, TDCP = True digestible crude protein, TDAA = True digestible amino acid.

* p<0.05; ** p<0.01.

Table 5. Stepwise regression with all variables or gross energy removed from the model (DM basis, n = 23)

TME _n equation ¹	Regression coefficient					Statistical parameter ²				
	Intercept	GE	EE	a*	b*	RMSE	R-square	BIC	C(p), Mallows	p-value
Equation 1	-2,911.3	1.04	-	-	-	199.3	0.65	249.3	-	**
p-value	**	**	-	-	-	-	-	-	-	-
Equation 2	-2,995.6	0.88	-	49.63	-	190.8	0.69	248.8	-2.87	**
p-value	**	**	-	**	-	-	-	-	-	-
Equation 3	992.6	-	-	-	43.89	259.5	0.40	260.2	-	**
p-value	*	-	-	-	**	-	-	-	-	-
Equation 4	NOINT	-	59.37	-	42.64	245.2	0.99	257.4	-	**
p-value	NOINT	-	**	-	**	-	-	-	-	-
Equation 5	NOINT	-	57.88	87.62	-	223.5	0.99	254.3	-4.43	**
p-value	NOINT	-	**	**	-	-	-	-	-	-

¹Equation 1 and 2 include all variables in the model; equation 3, 4 and 5 removed the gross energy from the model.

²RMSE = Root mean square error; R-square is the coefficient of determination and an adjusted R-square was calculated using the Noint option when the intercept was excluded from the model, p>0.15. BIC = Sawa's Bayesian Information Criteria; and C(p) is the Mallows statistic.

* p<0.05; ** p<0.01.

and Fiene et al. (2006), respectively, but similar to the results reported by Parsons et al. (2006) and Pahn et al. (2009). Many factors contribute to the large variability in chemical composition of DDGS, which include ethanol sources, fermentation technology (Cromwell et al., 1993), ratio of distillers grains to distillers solubles, drying temperatures and drying time (Noll et al., 2006). In the present study, DDGS samples were collected from 8 provinces of China, including northeast, northwest and central plains regions. Thus, corn varieties, climate and soil conditions in these areas may be a source of the differences.

The regression equation based on the whole data (n = 25) cannot predict TME_n precisely (R² = 0.34; RMSE = 361.9, data not shown), so the outliers were removed from the data set for further analysis. Because of the high correlation between GE and TME_n and the ratio of TME_n to GE being relatively stable, the outliers were removed when the ratio (TME_n to GE) was below or above 2 SD of the average value. Unlike the previous report, no proximate composition parameters were included in the model when gross energy and a* were in the equation. Rochell et al. (2011) reported that nutrients including NDF, GE and CP can be used to generate AME_n prediction equations for corn co-products (R² = 0.87), but they did not observe a correlation between AME_n and GE (r = 0.214, p = 0.44). Gross energy was strongly correlated with TME_n in the present study.

Gross energy usually cannot be determined by the feed industry in China; therefore, the present study wanted to develop a more practical model for commercial application with proximate composition and color scores. Fat and a* were in the final equation without an intercept. Similar to the results in this study, Batal and Dale (2006) found the best single predictor of TME_n was EE and followed by CF,

CP and ash, but these equations may only be used as a general guide (R² = 0.45). In contrast to the results of the current study, Cozannet et al. (2011) predicted AME_n of wheat DDGS with L* (R² = 0.77) instead of a*, while this study did not determined a* or b* values.

Results of TME_n showed considerable variation among samples (CV = 17.1%, n = 25) and the range spanned 1,610 kcal/kg. Small variation in TME_n values of DDGS, however, was reported by Parsons et al. (2006), who determined 20 DDGS on samples from feed mills in Minnesota with a CV of 3.6%; another study (Batal and Dale, 2006) determined 17 DDGS samples from 6 different plants in the Midwestern United States with the TME_n ranging from 2,490 to 3,190 kcal/kg (86% DM basis) and a CV of 6.4%. Although DDGS samples were high in fat, lower TME_n (Average = 2,144 kcal/kg, 46% of GE) was observed in the present study compared to the values in the United States in which TME_n takes nearly 60% of GE (Fastinger et al., 2006; Kim et al., 2010). One possibility of low TME_n is that DDGS in the present study were high in NDF, which cannot be digested by poultry (except turkey), and a high fiber diet enhanced intestinal movement that reduced feed retention time and feed digestibility. It is well known that most starch and sugar in DDGS were fermented to ethanol, and less than 7.0% starch is in the DDGS (Belyea et al., 2004; Stein et al., 2006). NFE is considered as a measure of the digestible carbohydrates (Leeson and Summers, 2009), thus another possible contributor to a low TME_n could be the low concentration of NFE in DDGS samples.

In contrast to previous reports, no correlation was found between color scores and TDAA or true digestible lysine. Previous studies demonstrated that sources of DDGS with a lighter and more yellow color tend to have greater TDAA and true digestible lysine (Batal and Dale, 2006; Babcock et

al., 2008; Pahn et al., 2009). Colors of DDGS are changed by the drying process (drying temperature and time), in part, this represents the degree of the Maillard-reaction, and this reaction decreased AA digestibility of DDGS. However, factors more than heat processing also affect the products color, darker and less yellow color was produced by increasing the soluble fraction into DDGS (Noll et al., 2005; Noll et al., 2007), but this change has no substantial effect on digestibility of AA. This suggests that color scores are not good indicator of AA digestibility for the present DDGS samples.

In addition, no correlation was found between CP and TAA in this study, and NAACP had a large variation between samples (CV = 55.01%). These results indicated that CP or color scores could not represent amino acid nutritional characteristics of DDGS.

In summary, the present results indicate that corn DDGS in China have a variable chemical composition and TME_n. In addition, gross energy is the best single indicator of TME_n. Finally, in this study, color score cannot predict TDAA or TD lysine of DDGS.

ACKNOWLEDGEMENTS

The work was supported by the Science & Technological Project of Shaanxi Province, China (2011KTDZ02-01-01, 2011KTCQ02-02), the National Key Technologies R&D Program of China (2011BAD26B03) and the R&D Fund of Shenzhen Kondarl Gaoling Feed Co. Ltd., Shaanxi, China.

REFERENCES

- AOAC International. 2006. Official methods of analysis of AOAC International. 18th ed. AOAC Int., Gaithersburg, MD, USA.
- Babcock, B. A., D. J. Hayes and J. D. Lawrence. 2008. Using distillers grains in the U.S. and international livestock and poultry industries. Midwest Agribusiness Trade Research and Information Center, Iowa State University, Ames, IA, USA.
- Batal, A. B. and N. M. Dale. 2006. True metabolizable energy and amino acid digestibility of distillers dried grains with solubles. *J. Appl. Poult. Res.* 15:89-93.
- Belyea, R. L., K. D. Rausch and M. E. Tumbleson. 2004. Composition of corn and distillers dried grains with solubles from dry grind ethanol processing. *Bioresour. Technol.* 94:293-298.
- Cozannet, P., M. Lessire, C. Gady, J. P. Métayer, Y. Primot, F. Skiba and J. Noblet. 2010. Energy value of wheat dried distillers grains with solubles in roosters, broilers, layers, and turkeys. *Poult. Sci.* 89:2230-2241.
- Cromwell, G. L., K. L. Herkelman and T. S. Stahly. 1993. Physical, chemical, and nutritional characteristics of distillers dried grains with solubles for chicks and pigs. *J. Anim. Sci.* 71:679-686.
- Ergul, T., C. M. Amezcua, C. Parsons, B. Walters, J. Brannon and S. Noll. 2003. Amino acid digestibility in corn distillers dried grains with solubles. *Poult. Sci.* 82(Suppl.1):70. (Abstr.)
- Farhat, A., L. Normand, E. R. Chavez and S. P. Touchburn. 1998. Nutrient digestibility in food waste ingredients for Pekin and Muscovy ducks. *Poult. Sci.* 77:1371-1376.
- Fastinger, N. D., J. D. Latshaw and D. C. Mahan. 2006. Amino acid availability and true metabolizable energy content of corn distillers dried grains with solubles in adult cecectomized roosters. *Poult. Sci.* 85:1212-1216.
- Fiene, S. P., T. W. York and C. Shasteen. 2006. Correlation of DDGS IDEA™ digestibility assay for poultry with cockerel true amino acid digestibility. *Proc. 4th Mid-Atlantic Nutr. Conf., Univ. Maryland, College Park.* pp. 82-89.
- Kim, E. J., C. M. Parsons, R. Srinivasan and V. Singh. 2010. Nutritional composition, nitrogen-corrected true metabolizable energy, and amino acid digestibilities of new corn distillers dried grains with solubles produced by new fractionation processes. *Poult. Sci.* 89:44-51.
- Leeson, S. and J. D. Summers. 2009. Commercial poultry nutrition. 3rd ed. Nottingham University Press, Thrumpton, Nottingham.
- National Research Council. 1994. Nutrient Requirements of Poultry. 9th rev, ed. Nalt. Acad. Press, Washington, DC, USA.
- Noll, S. L., J. Brannon and C. M. Parsons. 2007. Nutritional value of corn distiller dried grains with solubles (DDGs): Influence of solubles addition. *Poult. Sci.* 86(Suppl 1):68(Abstr.).
- Noll, S. L., C. M. Parsons and B. Walters. 2006. What's new since September 2005 in feeding distillers co-products to poultry. *Proc. Proceedings from the 67th Minnesota Nutrition Conference, University of Minnesota.* pp. 149-154.
- Pahn, A. A., C. S. Scherer, J. E. Pettigrew, D. H. Baker, C. M. Parsons and H. H. Stein. 2009. Standardized amino acid digestibility in cecectomized roosters and lysine bioavailability in chicks fed distillers dried grains with solubles. *Poult. Sci.* 88:571-578.
- Parsons, C. M. 1985. Influence of caeectomy on digestibility of amino acids by roosters fed distillers' dried grains with solubles. *J. Agric. Sci.* 104:469-472.
- Parsons, C. M., C. Martinez, V. Singh, S. Radhadkrishman and S. Noll. 2006. Nutritional value of conventional and modified DDGS for poultry. *Proc. Multi-State Poultry Nutrition and Feeding Conference.*
- Pedersen, C., M. G. Boersma and H. H. Stein. 2007. Digestibility of energy and phosphorus in ten samples of distillers dried grains with solubles fed to growing pigs. *J. Anim. Sci.* 85:1168-1176.
- Rochell, S. M., B. J. Kerr and W. A. Dozier. 2011. Energy determination of corn co-products fed to broiler chicks from 15 to 24 days of age, and use of composition analysis to predict nitrogen-corrected apparent metabolizable energy. *Poult. Sci.* 90:1999-2007.
- SAS Institute. 2003. Statistical Analysis Guide: Statistics. Ver. 8.1 ed. SAS Inst. Inc., Cary, NC, USA.
- Shim, M. Y., G. M. Pesti, R. I. Bakalli, P. B. Tillman and R. L. Payne. 2011. Evaluation of corn distillers dried grains with solubles as an alternative ingredient for broilers. *Poult. Sci.* 90:369-376.
- Spiehs, M. J., M. H. Whitney and G. C. Shurson. 2002. Nutrient

- database for distiller's dried grains with solubles produced from new ethanol plants in Minnesota and South Dakota. *J. Anim. Sci.* 80:2639-2645.
- Stein, H. H., M. L. Gibson, C. Pedersen and M. G. Boersma. 2006. Amino acid and energy digestibility in ten samples of distillers dried grain with solubles fed to growing pigs. *J. Anim. Sci.* 84:853-860.
- Van Soest, P. J., J. B. Robertson and B. A. Lewis. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74:3583-3597.
- Waldroup, P. W., Z. Wang, C. Coto, S. Cerrate and F. Yan. 2007. Development of a standardized nutrient matrix for corn distillers dried grains with solubles. *Int. J. Poult. Sci.* 6:478-483.